

***HST* and *UKIRT* Observations of the Center of the Trifid Nebula:  
Evidence for the Photoevaporation of a Proplyd and a  
Protostellar Condensation**

F. Yusef-Zadeh

Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208

(zadeh@northwestern.edu)

J. Biretta

STScI, 3700 San Martin Drive, Baltimore, MD 21218 (biretta@stsci.edu)

T. R. Geballe

Gemini Observatory, 670 N. A'ohoku Pl., Hilo, HI 96720 (tgeballe@gemini.edu)

Received \_\_\_\_\_; accepted \_\_\_\_\_

## ABSTRACT

The Trifid nebula (M20) is a well-known prominent optical HII region trisected by bands of obscuring dust lanes and excited by an O7.5 star HD 164492A. Previous near-IR, mid-IR and radio continuum observations of the cluster of stars at the center of the Trifid nebula indicated circumstellar disks associated with hot stars with envelopes that are photoionized externally by the UV radiation from the hot central star, HD 164492A. Using WFPC2 of the HST, we present evidence of a resolved proplyd in  $H\alpha$  and [SII] line emission from a stellar source emitting cool dust emission. Using UKIRT, an infrared observation of the stellar source with a proplyd indicates a late F – mid G spectral type. We also note a remarkable complex of filamentary and sheet-like structures which appear to arise from the edge of a protostellar condensation. These observations are consistent with a picture in which the bright massive star HD 164492A is responsible for the photoevaporation of protoplanetary disks of other less-massive members of the cluster as well as the closest protostellar condensation facing the central cluster. Using the evidence for a proplyd, we argue that the massive and intermediate mass members of the cluster, HD 164492C (B6 star) and HD 164492 (Herbig Be star) have disks associated with them.

*Subject headings:* ISM: HII regions—ISM: individual (M20)—ISM: planetary disks—Stars: formation

## 1. Introduction

Circumstellar disks are an integral part of star formation. Their existence has been established through a variety of millimeter, infrared and optical observations. HST observations have played an important role in transforming of our understanding of circumstellar material distributed within or close to an HII region powered by a cluster of massive stars. The so-called “proplyds”, or protoplanetary disks, and candidate proplyds have been found in the Orion Nebula (e.g., Laques & Vidal 1979; Garay et al. 1987; Churchwell et al. 1987; O’Dell 2001; Bally et al. 1998) in the Lagoon nebula (M8) (Stecklum et al. 1998), NGC 3603 (Brandner et al. 2000), Trifid nebula (M20) (Yusef-Zadeh, et al. 2000; Lefloch et al. 2002) and Carina nebula (Smith, Bally and Morse 2003). The Trifid nebula is a well-known HII region that is estimated to be  $\sim 10^5$  years old and star formation has taken place at the edge of the region within the last  $10^4$  years (Cernicharo et al 1998; Lefloch and Cernicharo 2000; Hester et al. 1999). It therefore presents us with an opportunity to examine proplyds at an stage of their exposure to the ionizing flux as well as to investigate the type of stars that proplyds can be associated with.

M20 is approximately  $10'$  in diameter with a distance estimated to be at the distance of 1.68 kpc (e.g., Lynds et al. 1985; see also Kohoutek et al. 1999) and is centered on a small cluster of hot stars dominated by components A through G of HD 164492. The ionizing flux required to maintain the HII region,  $10^{48.8} \text{ s}^{-1}$ , is supplied by the O7.5III star HD 164492A (e.g., Walborn 1973; Yusef-Zadeh et al. 2000), which has  $M_v = -5.3$  for  $A_v \approx 1.3$  towards the central stars (Lynds and O’Neil 1985). Several molecular condensations associated with protostellar sources lie within or at the edge of the HII region suggesting that a new generation of massive star formation has been induced by the expansion of the nebula (Cernicharo et al. 1998; Lefloch & Cernicharo 2000; Lefloch et al. 2002).

Radio and Infrared studies of the central region of M20 have suggested the presence of circumstellar disks in the vicinity of HD 164492A. In one study, three compact radio continuum sources with thermal spectrum was detected within  $0.''1$  of the cluster members identified by Gham et al. (1983) as a A2 IA star (HD 164492B), a B6 V star (HD 164492C) and D (Yusef-Zadeh

et al. 2000). HD 164492D lies only  $2''$  west of star C and is classified as a Herbig Be-star LkH $\alpha$  123 (Herbig 1957). The D star is also reported to be the brightest point source at 12.5 and 17.9  $\mu\text{m}$  images (Yusef-Zadeh et al. 2000). By analogy with M42, radio continuum emission from these sources was argued to be due to proplyds being photoevaporated by the intense UV field of HD 164492A (Yusef-Zadeh et al. 2000). The case for this idea proceeded along the lines developed by Garay et al. (1987) and Churchwell et al. (1987) for the proplyds first detected in radio continuum in M42. Another line of evidence for the existence of circumstellar disks came from near-IR L band observations (Yusef-Zadeh et al. 2000) showing several sources which were very dim or unseen shortward of K band image. Additional support for circumstellar disks surrounding the cluster members came from ISOCAM observations where several infrared sources were detected (Lefloch et al. 2002). These observations indicated two-temperature model of dust emission from mid-IR sources ranging between 150-200K and 500-1000K. The dust emission from HD 164492C and D implied the presence of warm neutral material around hot stars. Here, we present conclusive evidence for a proplyd from a star coincident with a mid-IR source IRS3 based on the *Hubble Space Telescope* (HST) observations of the bright central region of M20.

## 2. Observations

### 2.1. HST

WFPC2 observations (9104) of M20 were completed in June 2002 in order to make a mosaic image of this nebula, to study proper motion of HH 399 jet in the southeast region of the nebula and to search for proplyds in the central region of the nebula. Proper motion study of HH 399 was recently described by Yusef-Zadeh, Biretta and Wardle (2004). Deep narrow band observations with overlapping exposures were made to make a mosaic image of the nebula. A detailed account of the large scale mosaic image will be given elsewhere. Here, we concentrate on observations of the central region of the nebula which contains a cluster of stars and a bright ionized rim known to be embedded within the HII region.

To study the center of M20, a sequence of short and long exposures of the central cluster with two different orientations separated by  $160^\circ$  were carried out in  $H\alpha$  (F656N,  $4\times 2$  s,  $4\times 20$  s,  $2\times 200$  s), [SII] (F673N,  $2\times 300$  s), [OIII] (F502N,  $4\times 1.4$  s,  $4\times 14$  s,  $2\times 260$  s), F547M ( $2\times 40$  s), [NII] (F658,  $4\times 1.4$  s,  $4\times 14$  s) and [OI] (F631,  $2\times 1$  s,  $4\times 10$  s). We used a range of exposure times to insure non-saturated images for the brighter stars, and used dithering to improve PSF sampling and subtractions. The Planetary Camera (PC) chip was centered on the central cluster for all the above observations.

## 2.2. UKIRT

A  $1.4\text{--}2.5\ \mu\text{m}$  spectrum of the central star of HST1, as labeled in Figure 1, was obtained at the United Kingdom 3.8m Infrared telescope (UKIRT) on the night of 2004 Aug 29 (UT), using the facility spectrograph UIST (Ramsay Howat et al. 2000). The  $0.24''$  slit of UIST was oriented north-south and observations were made in the standard stare/nod mode. The total exposure time was 4 minutes. Hipparcos 86814 (F6V) was used for rough flux calibration and removal of telluric absorption lines.

## 3. Results

### 3.1. Proplyd

Figure 1 shows a grayscale PC image of  $H\alpha$  line emission from stars at the center of M20. The five bright cluster members HD 164492A to E stars at the center of the nebula are labeled in Figure 1. Two stars called HST1 and HST2 are located about  $16.86''$  and  $9''$  to the southwest of HD 164492A at  $\alpha, \delta(J2000) = 18^h 02^m 23^s.53, -23^\circ 01' 51.87''$ . HST1 shows extended  $H\alpha$  and [SII] line emission centered at  $\alpha, \delta(J2000) = 18^h 02^m 23^s.22, -23^\circ 01' 35.66''$ . A faint extended emission is also detected from HST2 centered at  $\alpha, \delta(J2000) = 18^h 02^m 23^s.07, -23^\circ 01' 45.45''$  but the reality of the extended component needs to be confirmed. Geometrical distortion has been removed before

the  $H\alpha$  positions of these stars were measured, and the absolute and relative positions of these stars are accurate to within  $0.5''$  and  $0.01''$ , respectively.

Figures 2a,b show contour and grayscale representations of a close-up  $H\alpha$  view of HST1, respectively. The morphology of the extended emission is reminiscent of proplyds noted in M42 where a crescent-shaped structure is pointing toward the central bright star HD 164492A. The proplyd structure coincides within  $0.22''$  and  $0.34''$  in right ascension and declination of a variable star centered at  $\alpha, \delta(J2000) = 18^h02^m23^s, -23^\circ01'36''$  V3791 with its brightness varying between 12.9 and 15.1 magnitudes (Kholopov 1998). The integrated  $H\alpha$  and [SII] fluxes from the proplyd are estimated to be  $4 \times 10^{-14}$  and  $2.2 \times 10^{-15}$  erg s $^{-1}$  cm $^{-2}$ , respectively. These estimates are made after the contribution from the central star and the background emission have been subtracted, but before any correction for extinction has been made. In order to investigate the reality of the extended emission and a dark ring associated with HST1, a brighter field star was selected in the vicinity of the central cluster. The comparison of the field star with HST1 showed that the number of counts from the crescent structure is three times higher than the brightness of the diffracted ring detected from the field star. We also measured the number of counts from the dark ring associated with HST1 and compared it with the background emission. We found the the brightness of the dark ring is about 2 counts pixel $^{-2}$  lower than expected from the diffraction of starlight and the background. The same crescent structure was noted in two different exposures with different orientations as described above. These measurements suggest strongly that both the crescent structure and the dark ring surrounding HST1 are real. The outermost and innermost radii of the crescent structure are  $0.36''$  and  $0.22''$  whereas the innermost radius of the dark ring is  $< 0.14''$ .

In order to identify the spectral type of the central star HST1, UKIRT observation of this star was carried out. Figure 3 shows the 2.0-2.5  $\mu$ m portion of the spectrum. The spectrum has been smoothed to a resolving power of  $\sim 600$ . The Br  $\gamma$  line in the calibration star was removed prior to ratioing; thus the Br  $\gamma$  absorption line in HST1 is real; it is the only clearly identified feature in the spectrum. Its strength, the presence of weaker Brackett series lines in the H portion of the

spectrum, and the lack of CO overtone band absorption at  $2.3 \mu\text{m}$  and longward suggest that the star's spectral type is late F - mid G. Thus, it is unlikely that the central star HST1 contributes a significant fraction of the ionizing radiation for the proplyd.

In spite of the short PC1 exposures, HD 164492B, C and D stars were too bright to bring out any faint extended emission arising from these stars. However, we noted that both stars C and D show multiple components. The C star showed two bright peaks in  $\text{H}\alpha$ , [NI] and [OIII] images separated by  $0.1''$  whereas the D star shows two bright peaks in  $\text{H}\alpha$  separated by  $0.1''$ , one of which had an [OIII] counterpart. The high-resolution images suggest that HD 164492B, C and D stars are binaries.

Based on HST and UKIRT observations, we assume that HST1 has a protoplanetary disk that is being heated and ionized by the FUV ( $6 \text{ eV} < h\nu < 13.6 \text{ eV}$ ) and EUV ( $h\nu > 13.6 \text{ eV}$ ) radiation from HD 164492A. Assuming a projected distance of  $2.8 \times 10^4 \text{ AU}$  ( $16.9''$  at the distance of  $1.68 \text{ kpc}$ ) between the central star and V3791, the Lyman continuum flux  $J_0$  is estimated to be  $2.8 \times 10^{12} \text{ photons cm}^{-2} \text{ s}^{-1}$ . The morphology of the proplyd shows that the radius of the ionization front  $R_i$  is subtending an angular distance of  $0.2''$  or  $\approx 340 \text{ AU}$ . The total Lyman continuum flux intercepted at the radius of the ionization front is expected to be  $\pi R_i^2 J_0 \approx 2.2 \times 10^{44} \text{ ph s}^{-1}$  assuming that there is no visual extinction between the ionizing source and the proplyd. The observed  $\text{H}\alpha$  flux is estimated to be  $1.5 \times 10^{43} \text{ ph s}^{-1}$  assuming  $A_v = 1.3$  magnitudes along the line of sight to the Trifid. A number of factors may contribute to a discrepancy of a factor of 15 between the expected flux intercepted by the protoplanetary disk and the extinction corrected  $\text{H}\alpha$  flux. The discrepancy could be partly due to the differential extinction toward the central star and the proplyd. Alternatively, this discrepancy could be due to a significant number of recombinations in the gas between the proplyd and the ionizing star. Considering that the proplyd lies in the bath of radiation and is closer to the ionization front of the HII region, it is plausible that the incident ionizing flux is different than that resulting from the  $r^{-2}$  geometrical dilution.

Using the emission measure  $E (\text{cm}^{-6} \text{ pc}) = 4.9 \times 10^{17} I_\alpha \approx n_e^2 L$  (see Bally and Reipurth 2001), the electron density  $n_e (\text{cm}^{-3})$  at the ionization front can be estimated from the observed

H $\alpha$  intensity  $I_\alpha \approx 2 \times 10^{-13}$  ergs s $^{-1}$ cm $^{-2}$ arcs $^{-2}$ . Assuming that L is the radius of the cusp  $\approx 2.5 \times 10^{-3}$  pc or 504 AU (0.3'' at the distance of 1.68 kpc),  $n_e \approx 1.1 \times 10^4$  cm $^{-3}$  after accounting for the extinction between us and M20. Given this emission measure, the estimated radio continuum flux density is estimated to be  $\approx 0.1$  mJy at 8GHz assuming the temperature of 8000  $^0$ K and the conversion factor used by Brandner et al. (2000). The total mass of ionized gas is estimated to be  $M_{HII} \approx 4/3 \pi \mu n_e R_i^3 \approx 2.1 \times 10^{-5} M_\odot$  where the mean molecular weight per electron  $\mu$  is assumed to be  $1.4 m_H$ . Estimating the expansion time to be  $R_i/(10\text{km/s}) \approx 160$  years, the mass-loss rate is estimated to be  $1.3 \times 10^{-7} M_\odot \text{yr}^{-1}$ .

### 3.2. Photoevaporative Ionized Rim

Evidence for the support of photoevaporative process being at work on a large scale comes from the striking distribution of the ionized gas  $\approx 25''$  away from HD 164492A. Figure 4 shows an HST mosaic image of the central region of M20. The mosaic image is constructed by combining H $\alpha$ , [SII] and [OIII] line images. Two bright ionized rims to the north and southeast delineate the surface of molecular condensations facing the central cluster. The so-called molecular condensation TC1 (the nomenclature used by Cernicharo et al. 1998) to the north is the closest protostellar condensation to the central cluster in M20. Figure 5a,b show a close-up view of the head of TC1 with a cometary morphology. A remarkable complex of elongated structures parallel and perpendicular to the ionized rim is best noted in H $\alpha$  and [OIII] line images. These features have no [SII] counterparts. The elongated structures extend up to  $12''$  or  $\sim 0.1$  pc, some of which appears to show a helical structure in projection and some sheet-like appearance. The lack of [SII] line emission from the filaments away from the ionized rim may be the result of an ionization stratification noted in the photoionized photoevaporative flow as seen in the HST image of M16 (e.g., Hester et al. 1996). Unlike the photoevaporative flows streaming normal to the surface of the dark cloud toward the ionizing star, the features noted in Figures 4 and 5 make a large angle with respect to normal to the surface of the molecular condensation.



#### 4. Discussion

The evidence for the existence of protoplanetary disks at the center of M20 being illuminated by a bright O7.5 star HD 164492A comes from a number of direct and indirect measurements. The most conclusive evidence comes from the optical morphology of  $H\alpha$  and [SII] line emission arising from a crescent-shape structure with its cusp pointing toward HD 164492A star. The proplyd coincides with a late F – mid G spectral type and an infrared source IRS 3 which shows evidence of warm dust emission from the protoplanetary disk. The physical size or the cusp radius of the proplyd is  $\approx 8 \times 10^{15}$  cm which is larger than the typical size of the proplyds found in the Orion nebula. Radio continuum, near-IR and mid-IR measurements suggest dense material in the form of protoplanetary disks being photoevaporated by HD 164492A. These stars include a massive star HD 164492C or IRS 1 and an intermediate mass star HD 164492D or IRS 2. The evidence of proplyd from IRS 3 provides indirect support for the hypotheses that the disks of these massive stars IRS 1 and IRS 2 are also photoevaporated by HD 164492A but because of their high stellar brightness, their corresponding proplyds were not detected.

One of the key questions is whether or not proplyds survive sufficiently long for planet formation to occur (Churchwell et al. 1987, Henney & O'Dell 1999). To examine the properties of the proplyd in M20, we estimate the lifetime of the neutral disk using the column density  $N(\text{HI}) \approx 10^{21} \text{cm}^{-2}$  estimated from mid-IR observations (Lefloch et al. 2002). These authors derive a typical column density of hydrogen for the cool component of the dust emission from IRS 1, 2 and 3, all of which show similar spectral characteristics. If we assume that this cool column of gas comes from a disk of radius  $R_H \approx 0.14''$  or 235 AU, the total mass of the disk  $M_H \approx N_H \pi R_H^2 m_{H2} \approx 8.8 \times 10^{-5} M_\odot$ . The mass of cold neutral material is estimated to be similar to the mass of ionized gas, thus the emission is argued to be originating from the photon-dominated region PDR of a reservoir of dense circumstellar material exposed to the ionizing radiation of HD 164492A (Yusef-Zadeh *et al.* 2000; Lefloch *et al.* 2001). The lifetime of the PDR in the proplyd associated with IRS 3 is estimated to be roughly 500 years. However, this timescale is clearly underestimated since the column density of molecular gas is unknown.

Additional evidence for the support of photoevaporative process being at work on a large scale comes from the distribution of a bright rim seen  $\approx 36''$  away from HD 164492A. The bright ionized rim delineates the surface of a molecular condensation (TC1) facing the central cluster. Figures 4, 5 show a remarkable extended structure projected against the surface of the ionized rim as best noted in  $H\alpha$  and [OIII] lines. This shock-like filamentary structures with size scale of about 0.1 pc are detected only at the edge of the ionization front where the incident radiation from HD 164492A is non-normal to the surface of the molecular condensation. We speculate that the origin of such a structure could be due to a number of proplyd outflows distributed in the vicinity of TC1. The nature of several stars projected against the filamentary structure needs further investigation to determine their relationship to the filamentary structure. In this picture the proplyd outflows could be shocked by their interaction with each other and/or with the stellar wind from HD 164492A. Similar processes are known to take place in the Orion nebula (e.g., Garcia-Arrendono, Henny & Arthur 2001; Henney 2002; Graham et al. 2002). Other effects that could contribute to the production of filamentary [OIII] structure at such curious location with respect to the central HD 164492A star is the onset of instability of the ionization front. Recent numerical work suggests that the ionization front can become unstable when the source of radiation makes an inclined angle with respect to the surface of TC1 (e.g., Williams 2002). Obviously, more detailed work is necessary to understand the nature of these diffuse ionized features.

**Acknowledgments:** We wish to thank R. O'Dell and Mark Wardle who were co-investigators on the HST proposal under which these observations were carried out. We are also grateful to the referee, William Henney whose comments improved the manuscript. In particular, his suggestions to account for the origin of the large-scale diffuse features in the vicinity of the protostellar condensation were extremely helpful. TRG thanks the staff of UKIRT; his research is supported by the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy on behalf of the international Gemini partnership of Argentina, Australia, Brazil, Canada, Chile, the United Kingdom, and the United States of America.

## REFERENCES

- Bally, J. & Reipurth, B. 2001, ApJ, 546, 299
- Bally, J., Sutherland, R.S., Devine, D. & Johnstone, D. 2000, ApJ, 116, 293
- Brandner, W., Grebel, E.K., Chu, Y.H., Dottori, H., Brandl, B., Richling, S., York, H.W., Points, S.D., Zinnecker, H., 2000, AJ, 119, 292
- Cernicharo, J., Lefloch, B., Cox, P., Cesarsky, D., Esteban, C., Yusef-Zadeh, F., Mendez, D.I., Acosta-Pulido, J., Lopez, R. J. Garcia, & Heras, A. 1998, Science, 282, 462
- Churchwell, E., Felli, M., Wood, D.O.S. & Massi, M. 1987, ApJ, 321, 516
- Gahm, G.F., Ahlin, P. and Lindroos, K.P. 1983, A&AS 51, 143
- Garay, G., Moran, J. M. & Reid, M. J. 1987, ApJ, 314, 535
- Garcia-Arrendondo, F., Henney, W.J. & Arthur, S.J. 2001, ApJ, 561, 830
- Graham, M.F., Meaburn, J., Garrington, S.T., O’Brien, T.J., Henney, W.J. & O’Dell, C.R. 2002, ApJ, 570, 222
- Henney, W.J. 2002, RMxAAm 38, 71
- Herbig, G. H. 1957, ApJ., 125, 654
- Henney, W.J. & O’Dell, C.R. 1999, AJ, 118, 2350
- Hester, J., Scowen, P.A., Sankrit, R., Lauer, T.R., Ajhar, E.A. et al. 1996, AJ, 111, 2349
- Hester, J., Scowen, P., Stappelfeldt, K. & Krist, J. 1999, BAAS, 194, 68.10
- Kholopov P.N., Samus, N.N., Frolov, M.S., Goranskij V.P., Gorynya, N.A. et al. 1998 in Combined General Catalogue of Variable Stars, 4.1
- Kohoutek, L., Mayer, P. & Lorenz, R. 1999, A&AS, 134, 129
- Laques, P. & Vidal, J.L. 1979, A&A, 73, 97
- Lefloch, B. & Cernicharo, J. 2000, ApJ, 545, 340

- Lefloch, B., Cernicharo, J., Rodriguez, L.F., Miville-Deschenes, M.A., Cesarski, D. & Heras, A. 2002, *ApJ*, 581, 335
- Lynds, B.T. and O’Neil, E.J. Jr. 1985, *ApJ*, 294, 578
- Lynds, B.T., Canzian, B.J. & O’Neil, E.J., Jr. 1985, *ApJ*, 288, 164
- O’Dell, C. R. 2001, *ARA&A*, 39, 99
- Panagia, N. 1973, *AJ*, 78, 929
- Ramsay Howat, S. K., Ellis, M.A., Gostick, D.C., Hastings, P. R., Strachan, M., & Wells, M. 2000, *Proc. SPIE*, 2198, 350
- Stecklum, B., Henning, T., Feldt, M., Hoare, M.G., Hofner, P., Richer, R. 1998, *AJ*, 115, 767
- Smith, N., Bally, J. and Morse, J.A. 2003, *ApJ*, 587, L105
- Spitzer, L. 1978, “Physical Processes in the Interstellar Medium” (New York: Interscience)
- Williams, R.J.R. 2002, *MNRAS*, 331, 693
- Walborn, N.R. 1973, *AJ*, 78, 1067
- Yusef-Zadeh, F., Biretta, J. & Wardle, M. 2005, *ApJ*, 624, 246
- Yusef-Zadeh, F., Shure, M., Wardle, M. & Kassim, N. 2000, *ApJ*, 540, 849

---

This manuscript was prepared with the AAS L<sup>A</sup>T<sub>E</sub>X macros v4.0.

Fig. 1.— PC1 image of H $\alpha$  emission from the central cluster of stars in M20 based on  $2 \times 200$  sec exposures. The reference pixel position is at  $\alpha, \delta(J2000) = 18^h 02^m 23^s.2, -23^\circ 01' 38''$ . The labeled stars A to E correspond to cluster members HD 174492A – E.

Fig. 2.— (a) Grayscale representation of the  $H\alpha$  distribution of HST1 using PC1 [left panel]. (b) Contours of  $H\alpha$  emission from the protostellar disk with levels at (18, 20, 22, 24, 26, 28 and 30)  $\times 2.051 \times 10^{-17} \text{ergcm}^{-2}\text{s}^{-1}$

Fig. 3.— The 2.0-2.5  $\mu\text{m}$  spectrum of the central star of HST1, shown at  $R \sim 600$ . The location of the Br  $\gamma$  line is indicated.

Fig. 4.—  $H\alpha$ , [SII] and [OIII] images of the inner  $3'$  of M20. Two bright ionized rims associated with two molecular condensations are noted to the north and southeast of the stellar cluster at the center of the nebula.

Fig. 5.— (a)  $H\alpha$  (a) and [OIII] (b) images of the northern region of the bright ionized rim of the Trifid nebula where filamentary structures are detected.

This figure "f1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>

This figure "f2a.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>

This figure "f2b.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>



This figure "f3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>

This figure "f4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>

This figure "f5a.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>

This figure "f5b.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0505155v1>